

Constraining the inflaton potential with gravitational waves from oscillons

JAN TRÄNKLE (3rd YEAR PHD @ ITP HANNOVER U.)

INFLATION 2025, IAP, PARIS, 05.12.2025

BASED ON: 2512.XXXXX WITH K. LOZANOV & M. SASAKI



Motivation

- unknown expansion history between end of inflation and radiation domination





Figure: Allahverdi et al. [2006.16182]



Constraining the **inflaton potential** with gravitational waves from oscillons

Models

- Inflaton potentials favoured by CMB data




Figure: *Planck* 2018

Models

- α -attractor T-model

$$V(\phi) = \frac{m^2 M^2}{2} \tanh^{2n} \left(\left(\frac{|\phi|}{M} \right)^{1/n} \right)$$

Kallosch, Linde (2013)

- generalized axion monodromy

$$V(\phi) = \frac{m^2 M^2}{q} \left(\left(1 + \left(\frac{\phi}{M} \right)^2 \right)^{q/2} - 1 \right)$$

Silverstein, Westphal (2008)



- hilltop (squared)

$$V(\phi) = \frac{m^2 M^2}{2p^2} \left(1 - \left(\frac{\phi}{M} \right)^p \right)^2$$

Boubekeur, Lyth (2005)



Models

- Inflaton potentials favoured by CMB data



Models

- Inflaton potentials favoured by CMB data $\rightarrow$ supporting oscillons



Inflaton self-interaction

- expansion around quadratic minimum

$$V(\phi) = \frac{1}{2}m^2\phi^2 + V_{\text{nl}}(\phi)$$

$$V_{\text{nl}}(\phi \ll M) \approx g\phi^3 - \lambda\phi^4 + \dots$$

$$\lambda = c_\lambda \frac{m^2}{M^2}$$

$$g = c_g \frac{m^2}{M}$$

Amin et al. [1106.3335]

Amin [1006.3075]

Inflaton self-interaction

- expansion around quadratic minimum

$$V(\phi) = \frac{1}{2}m^2\phi^2 + V_{\text{nl}}(\phi)$$

$$V_{\text{nl}}(\phi \ll M) \approx g\phi^3 - \lambda\phi^4 + \dots$$

$$\lambda = c_\lambda \frac{m^2}{M^2}$$

$$g = c_g \frac{m^2}{M}$$

- attractive self-interaction $V_{\text{nl}}(\phi) < 0 \iff \lambda > 0$
- fragmentation for $M \lesssim 5 \times 10^{-2} M_{\text{Pl}}$

Amin et al. [1106.3335]

Amin [1006.3075]

Constraining the inflaton potential with gravitational waves from **oscillons**

Oscillon formation




Figure: Amin, Mocz [1902.07261],
Allahverdi et al. [2006.16182]

[Talk by Josu Aurrekoetxea]



Oscillon formation




Figure: Amin, Mocz [1902.07261],
Allahverdi et al. [2006.16182]

Oscillon properties

- oscillon profile: oscillating core + small radiation tail




Figure: Zhang et al. [2004.01202]

Oscillon properties

- oscillon profile: oscillating core + small radiation tail
- remarkable lifetimes $\sim (10^6 - 10^8)m^{-1}$
- single oscillon mass $m_{\text{osc}} = \mathcal{O}[10^2] \frac{M^2}{m} \gg m$





Figure: Zhang et al. [2004.01202]

Oscillon properties

- oscillon profile: oscillating core + small radiation tail
- remarkable lifetimes $\sim (10^6 - 10^8)m^{-1}$
- single oscillon mass $m_{\text{osc}} = \mathcal{O}[10^2] \frac{M^2}{m} \gg m$
- rapid final decay to (scalar) radiation




→ enhanced production of induced GWs !

Inomata et al. [1904.12879]

Constraining the inflaton
potential with **gravitational
waves** from oscillons

Oscillon induced GWs



mean separation $\sim k_{\text{osc}}^{-1}$


Poissonian number density fluctuations δ_{osc}

$$\mathcal{P}_{\delta_{\text{osc},f}} \propto (k/k_{\text{osc}})^3$$

Figure: Amin, Mocz [1902.07261]

G. Domènech [2109.01398]

Oscillon induced GWs



mean separation $\sim k_{\text{osc}}^{-1}$

Poissonian number density fluctuations δ_{osc}


$$\mathcal{P}_{\delta_{\text{osc},f}} \propto (k/k_{\text{osc}})^3$$

$$\text{⚙} G_{\mu\nu}^{(1)} = T_{\mu\nu}^{(1)} \Rightarrow \text{curvature perturbation } \Phi$$

Figure: Amin, Mocz [1902.07261]

G. Domènech [2109.01398]

Oscillon induced GWs



mean separation $\sim k_{\text{osc}}^{-1}$

Poissonian number density fluctuations δ_{osc}

$$\mathcal{P}_{\delta_{\text{osc},f}} \propto (k/k_{\text{osc}})^3$$


⚙ $G_{\mu\nu}^{(1)} = T_{\mu\nu}^{(1)} \Rightarrow$ curvature perturbation Φ

⚙ $G_{\mu\nu}^{(2)} = T_{\mu\nu}^{(2)} \Rightarrow$ scalar induced GWs h_{ij}

Figure: Amin, Mocz [1902.07261]


G. Domènech [2109.01398]

Induced GW spectrum



$$\beta \equiv \frac{\rho_{\text{osc},f}}{\rho_f} = 0.8$$

Induced GW spectrum




Constraining the inflaton potential with gravitational waves from oscillons

Parameter space

$$\lambda = c_\lambda \frac{m^2}{M^2}$$

fixed A_s, n_s
from *Planck*




Parameter space

$$\lambda = c_\lambda \frac{m^2}{M^2}$$

fixed A_s, n_s
from *Planck*


$$r < 10^{-3}$$

$$r < 0.056$$





Parameter space

$$g = c_g \frac{m^2}{M}$$



Parameter space

- *caveats*: errorbars from simulations (lifetimes, mass, ...), additional GW sources, ...



Conclusion

- oscillon formation & domination generically expected in many models
- rapid final decay of oscillons leads to loud induced GW signal
- GW overproduction provides new constraints on parameter space


Conclusion

- oscillon formation & domination generically expected in many models
- rapid final decay of oscillons leads to loud induced GW signal
- GW overproduction provides new constraints on parameter space



Thank you for your attention!

... Questions ?



Additional Material

Relating parameters & observables

- potential slow-roll parameters

$$\epsilon_V(\phi_e) = \frac{M_{\text{Pl}}^2}{2} \left(\frac{V_\phi(\phi)}{V(\phi)} \right)^2 \Big|_{\phi_e} \simeq 1$$



$$\phi_e(M)$$




$$N_*(M)$$

$$n_s = 1 - 6\epsilon_V(\phi_*) + 2\eta_V(\phi_*)$$



$$\phi_*(M)$$

$$\eta_V = M_{\text{Pl}}^2 \frac{V_{\phi\phi}(\phi)}{V(\phi)}$$



$$A_s = \frac{1}{8\pi^2 \epsilon_V(\phi_*)} \frac{H_*^2}{M_{\text{Pl}}^2}$$



$$m(M)$$

$$A_s \simeq 2.1 \times 10^{-9}$$

$$n_s \simeq 0.965$$

$$r = 16\epsilon_V(\phi_*) = \frac{2}{A_s \pi^2} \frac{H_*^2}{M_{\text{Pl}}^2}$$



$$M < M_{\text{max}}$$

$$r < 0.056$$

[1] Planck 2018

(Classical) Decay rate

- decay rate (α -attractor T-model):
 3ω radiation suppressed in dip
→ quasi-stable attractor solution with
unique oscillon mass m_{osc}
→ extremely long lifetime





Figure: Zhang et al. [2004.01202]

Model	c_g	c_λ	$r < 0.056$	$r < 10^{-3}$	$\tau_{\text{osc}} = t_{\text{osc}} m$	N_{osc}	N_*	$\bar{H}_f$
α -attractor T ($n = 1$)	0	1/3	$M < 12M_{\text{Pl}}$	$M < 1.3M_{\text{Pl}}$	$\gtrsim 3 \times 10^6$	7	57	$1/\sqrt{6}$
Axion monodromy ($q = 1$)	0	1/8	$M \lesssim M_{\text{Pl}}^{\text{a}}$	(ruled out)	$\sim 3 \times 10^8$	11	71 ^a	$\sqrt{\frac{M_{\text{Pl}}}{3\sqrt{2}M}}$
Logarithmic ($q = 0$)	0	1/4	$M < 3.2M_{\text{Pl}}$	$M \ll M_{\text{Pl}}$	$\sim \mathcal{O}(1) \times 10^8$	9	29 – 44	0.415
Plateau ($q = -1$)	0	3/8	$M < 8.3M_{\text{Pl}}$	$M < 0.018M_{\text{Pl}}$	$\gtrsim 6 \times 10^8$	10	38 – 50	$1/\sqrt{3}$
Hilltop ($p = 4$)	3/2	-17/8	$M < 36M_{\text{Pl}}$	$M < 8.5M_{\text{Pl}}$	$\sim \mathcal{O}(1) \times 10^5$	4	54 – 86	$1/(4\sqrt{2})$
Hilltop ($p = 6$)	5/2	-155/24	$M < 51M_{\text{Pl}}$	$M < 8.5M_{\text{Pl}}$	$\sim \mathcal{O}(1) \times 10^6$	5	53 – 71	$1/(6\sqrt{2})$
Hilltop ($p = 8$)	7/2	-105/8	$M < 67M_{\text{Pl}}$	$M < 9.3M_{\text{Pl}}$	$\sim 2 \times 10^6$	5	52 – 67	$1/(8\sqrt{2})$

Simulations: Zhang et al. [2004.01202], Lozanov, Amin [1710.06851], Olle et al. [1906.06352], Antusch et al. [1907.00611]

Curvature perturbation

- isocurvature initial conditions $\Phi(0) = 0$



$$\propto \mathcal{C}_1(\beta) \delta_{\text{osc},f}$$

$$\propto \mathcal{C}_2(\beta) \delta_{\text{osc},f} \kappa^{-2} \quad \kappa = \left(\frac{k}{k_f} \right)$$

Poisson eq.
 $k^2 \Phi \propto \delta_{\text{osc}}$

$$\beta \equiv \frac{\rho_{\text{osc},f}}{\rho_f} = 0.8$$

Induced Gravitational Waves

- Perturbed FLRW metric

$$ds^2 = a^2(\tau) \left[-(1 - 2\Phi)d\tau^2 + (\delta_{ij} + 2\Phi\delta_{ij} + h_{ij}) dx^i dx^j \right]$$


$$G_{\mu\nu} = T_{\mu\nu}$$

- Tensor perturbations at 2nd order

$$h_k''(\tau) + 2\mathcal{H}h_k'(\tau) + k^2h_k(\tau) = \mathcal{S}_k(\tau) \propto (\Phi + \Phi'/\mathcal{H})^2$$

G. Domènech [2109.01398]

Induced Gravitational Waves

- Tensor perturbations at 2nd order

$$h_k''(\tau) + 2\mathcal{H}h_k'(\tau) + k^2 h_k(\tau) \propto (\Phi + \Phi'/\mathcal{H})^2$$

- Spectral GW energy density

$$\Omega_{\text{GW}}(k) \sim \langle h_{ij} h_{ij} \rangle \sim \int dv \int du \overline{I^2}(k\tau, u, v) \langle \Phi \Phi \rangle(uk) \langle \Phi \Phi \rangle(vk)$$

- largest contribution to GWs from rapid transition $w = 0 \rightarrow 1/3$ right after
oscillon decay (Poltergeist mechanism)

G. Domènech [2109.01398]

Inomata et al. [1904.12879]